1 (Sem-4) PHY 4

## 2025

## PHYSICS.

Paper: PHY0400404

(Mathematical Physics)

Full Marks: 45

Time: Two hours

The figures in the margin indicate full marks for the questions.

1. Answer to the following questions:

1×5=5

- (a) Express the Laplace equation for twodimensional Cartesian system.
- (b) What is the necessary condition for the existence of the Fourier's series?

- (c) If certain complex function f(z) is analytic then which equation should it satisfy?
- (d) What is the residue of a function at a simple pole?
- (e) What is the fundamental property of Levi-Civita symbol,  $\varepsilon_{ijk}$ ?
- 2. Answer any five questions:  $2 \times 5 = 10$ 
  - (a) Express the general form of Laplace equation in spherical polar coordinate system.
  - (b) State the Dirichlet conditions for the existence of a Fourier series.
  - (c) State the Cauchy-Riemann conditions in Cartesian coordinates.
  - (d) Express the difference between a pole and a branch point with examples.

- (e) Define a symmetric tensor with an example.
- (f) What is the significance of Einstein's summation convention?
- (g) Write down the probability mass function of the Poisson distribution and illustrate the parameters within.
- (h) Find the mean and variance of a binomial distribution with parameters n and p.
- (i) Express the Fourier series of a function f(x) = x defined within the bound  $(-\pi, \pi)$ .
- (j) What type of boundary conditions are used to solve the wave equation for a vibrating string fixed at both ends?

- 3. Answer any four questions:  $5 \times 4 = 20$ 
  - (a) Solve the one-dimensional wave equation for a string of length L fixed at both ends using the separation of variables method (Give the general form of the solution and mention the boundary conditions).
  - (b) Solve the Laplace's equation in two dimensions for a rectangular region with suitable boundary conditions using separation of variables method (express the solution graphically without evaluating the arbitrary constants).
  - Find the Fourier series (sine and cosine form) for the function  $f(x) = x^2$  in the interval  $(-\pi, \pi)$ . Show all steps clearly.
  - Expand the periodic square wave function in a complex Fourier series and write down the expression.

- State and prove Cauchy's integral formula for a function analytic in a simply connected domain. (State all assumptions clearly)
- Determine the nature and the order of the singularity of the function,

$$f(z) = \frac{\sin z}{z^3}$$
 at  $z = 0$ , and compute its residue.

- If  $A^{\mu\nu}$  and  $B_{\mu\nu}$  are two tensors of rank two, then describe their transformation rules under coordinate transformation and also show that  $A^{\mu\nu}B_{\mu\nu}$  is an invariant quantity.
- Starting from the Poisson distribution, derive the condition under which it approximates the binomial distribution. Explain the physical or statistical significance of this limit.

B06FN 0153

- 4. Answer **any one** question:  $10 \times 1 = 10$ 
  - (a) Solve the following differential equation using separation of variable method.

$$\frac{\partial u}{\partial x} = 2\frac{\partial u}{\partial t} + u$$
, where  $u(x, 0) = 6e^{-3x}$ .

(b) Given,  $f(x) = \begin{cases} -1, & \text{for } -\pi < x < -\frac{\pi}{2}, \\ 0, & \text{for } -\frac{\pi}{2} < x < \frac{\pi}{2}, \\ +1, & \text{for } \frac{\pi}{2} < x < \pi \end{cases}$ 

find the Fourier expression of f(x).

(c) Evaluate the integral  $\oint_C \frac{e^z}{z^2(z-1)} dz$ , where C is the positively oriented circle |z| = 2, using residue theorem. Clearly identify the singularities of the integrand, determine the order of each singularity, compute the residues at the poles enclosed by C. 6+4=10

6

(d) Prove the quotient law of tensors. Show that the Kronecker delta  $\delta^i_j$  behaves as a mixed tensor. What is its rank? Using tensor transformation laws, show that it acts as the identity operator under index contraction. 4+3+1+2=10